

Toxic Footprint: Exploring the Impact of E-Waste on Biodiversity and Ecosystems

¹Rahima Khatun*, ²Amrik Bhattacharya, ³Sayantika Mukherjee and ¹Amitlal Bhattacharya

¹Dukhulal Nibaran Chandra College, Aurangabad, Murshidabad, 742201

²Lakhuria High School, Kaliaganj, Nadia, West Bengal, 741150

³Amity University, Kolkata, West Bengal, 700135

*Corresponding author

Abstract:

The rapid advancement of technology has led to an unprecedented rise in electronic waste (e-waste), posing a serious threat to biodiversity and ecological balance. E-waste contains hazardous substances such as lead, mercury, cadmium, and brominated flame retardants, which can contaminate soil, water, and air when improperly disposed of. This paper examines the relationship between e-waste generation and biodiversity loss, highlighting the pathways through which toxic materials disrupt ecosystems. It also evaluates current recycling practices, identifies gaps in waste management systems, and explores sustainable solutions to mitigate environmental damage. The study emphasizes the urgent need for improved policies, public awareness, and environmentally responsible recycling methods to protect biodiversity and ensure ecological sustainability.

Key Words: E-waste, Biodiversity, Environmental Pollution, Recycling, Toxic Substances, Sustainability

Date of Submission: 03-04-2026

Date of acceptance: 14-04-2026

I. Introduction

The rapid expansion of digital technologies has transformed modern society, but it has also generated one of the fastest-growing waste streams: electronic waste (e-waste). According to the Global E-waste Monitor 2024, approximately 62 million tons of e-waste were produced in 2022, with projections reaching 82 million tons by 2030. However, recycling rates remain significantly lower than generation, creating a widening gap that poses serious environmental risks. Beyond resource loss, e-waste represents a major ecological threat due to the release of hazardous substances into natural systems.

E-waste is uniquely problematic because of its complex composition. Electronic devices contain valuable materials such as copper, gold, silver, and rare earth elements, alongside toxic substances including lead, mercury, cadmium, chromium, and brominated flame retardants. When improperly managed—through landfilling, open burning, or informal recycling—these pollutants are released via leaching, volatilization, and runoff. As a result, contaminants accumulate in soils, water bodies, and the atmosphere, disrupting environmental chemistry and biological systems.

The ecological consequences extend directly to biodiversity and ecosystem stability. Soil microorganisms, which are essential for nutrient cycling, are highly sensitive to heavy-metal contamination, and their decline can impair soil fertility and ecosystem productivity. In aquatic environments, toxic elements and persistent organic pollutants interfere with reproduction, development, and metabolism in aquatic organisms. Furthermore, bioaccumulation and biomagnification increase contaminant concentrations at higher trophic levels, affecting fish, birds, and mammals. These processes lead to reduced survival rates, reproductive failure, genetic damage, and behavioural changes, ultimately contributing to biodiversity loss and weakened ecosystem resilience.

The situation is exacerbated by the inefficiency of global recycling systems. In 2022, only about 22.3% of e-waste was formally collected and recycled, leaving a substantial portion unmanaged. Informal recycling practices, common in many developing regions, often involve hazardous methods such as open burning and acid leaching, which release toxins without proper containment. These activities frequently occur near populated or agriculturally significant areas, allowing pollutants to spread into ecosystems and food chains, thereby threatening both biodiversity and human health.

The growing urgency of the e-waste crisis is further driven by shortening product lifecycles and limited repair or reuse practices. Increasing demand for electronic devices, coupled with insufficient collection

infrastructure, has created a system where waste generation far exceeds recovery. This imbalance is particularly concerning in biodiversity-rich regions, where even low levels of contamination can have disproportionate ecological impacts.

This review explores the relationship between e-waste and biodiversity loss, focusing on contamination pathways, ecosystem-level effects, and current management limitations. It also discusses emerging solutions, including improved recycling technologies, regulatory frameworks, and circular design strategies, emphasizing the need to address e-waste as both an environmental and biodiversity challenge.

II. Composition and Characteristics of E-Waste

E-waste is a highly heterogeneous stream that contains both economically valuable resources and environmentally hazardous constituents. Its composition varies by product category and generation, but it typically includes metals, polymers, glass, ceramics, rare earth elements, and a range of additives and functional chemicals used in circuitry, displays, batteries, and housings. This dual character is what makes e-waste simultaneously a resource reservoir and a pollution source, especially when it is dismantled, burned, or dumped without adequate control.

2.1 Valuable Materials

A major reason e-waste has become central to the circular economy is that it contains high-value secondary raw materials that can be recovered and reused. Printed circuit boards, connectors, wiring, displays, and other components are rich in precious and base metals such as gold, silver, copper, and palladium, while some streams also contain platinum-group metals and rare earth elements used in magnets, screens, and phosphors. In many device categories, these metals make electronic scrap more resource-dense than natural ores, which is why e-waste is increasingly described as a form of “urban mining”. Beyond metals, e-waste also contains reusable polymers, glass, and composite materials that can be recovered if dismantling and separation are performed efficiently. However, the recovery potential is often undermined by product miniaturization, multi-material assemblies, and the presence of bonded or laminated layers that are difficult to separate mechanically. Recent analyses emphasize that the economic value of e-waste is substantial, yet global recycling efficiency remains low, meaning that large quantities of recoverable materials are still lost to informal disposal or low-yield processing routes.

2.2 Hazardous Substances

The environmental concern surrounding e-waste arises mainly from the toxic additives and heavy metals embedded in its components. Lead remains one of the most important contaminants, particularly in solder, cathode ray tube glass, batteries, and printed wiring boards, where it can affect nervous systems and contaminate soil and dust when released. Mercury is found in certain lighting units, switches, and display-related components, and it is especially dangerous because of its toxicity to aquatic organisms and its ability to move through food webs after environmental release. Cadmium is another major pollutant, commonly associated with rechargeable batteries, pigments, semiconductors, and some older electronic parts. It can damage kidneys and accumulate in soils and sediments, where it may persist for long periods and disrupt microbial and plant communities. In addition, brominated flame retardants and related halogenated compounds are widely used in plastic housings, circuit boards, and cable insulation to reduce flammability, but they are persistent organic pollutants that can generate toxic by-products during uncontrolled burning or thermal treatment. Recent reports also note that informal recycling and open burning can release a much broader mixture of contaminants than is often appreciated, including dioxins, furans, polycyclic aromatic hydrocarbons, and other combustion-derived toxins. These substances do not remain confined to the site of disposal; they can enter the atmosphere as smoke and particulates, infiltrate soil through ash deposition, and move into waterways through runoff and leaching. Once released, they can trigger chronic ecological damage by altering nutrient cycles, reducing species fitness, and increasing exposure risks across multiple trophic levels. The combination of valuable and hazardous materials makes e-waste a uniquely challenging waste stream. On one hand, it offers a significant source of secondary metals and reusable resources; on the other, it contains chemical hazards that can destabilize ecosystems if managed improperly. This is why updated reviews increasingly frame e-waste not simply as a disposal issue, but as a materials-management, pollution-control, and biodiversity-conservation problem at the same time. In practice, the environmental footprint of e-waste depends on how effectively it is collected, sorted, and processed. Where formal recycling systems are weak, hazardous substances are more likely to be dispersed into the environment, while valuable materials are lost from the resource cycle. For this reason, understanding the composition of e-waste is essential for designing safer recycling technologies, stronger regulations, and more sustainable product architectures.

III. Impact of E-Waste on Biodiversity

E-waste affects biodiversity mainly through the release of toxic metals and persistent organic pollutants into soil, water, and air, where they alter habitat quality and biological function.

Table 1. Diseases caused by E-waste in human beings.

Hazardous Components	Present In	Consequences of Hazardous Components in E-waste
Arsenic	Semiconductors, diodes, microwaves, LEDs, solar cells	The nervous system and the skin may be impacted. Long-term exposure can cause lung cancer.
Asbestos	Insulators in heating equipment	Breathing problems, coughing, lung damage, and even cancer are serious adverse effects.
Barium	Fillers for plastics and rubbers, as well as electron tubes	Heart muscle can be affected.
BFR	Different casing, circuit boards, chips	The reproductive and immune systems may be harmed. Hormone imbalances and endocrine system issues are possible.
CD	PCB, batteries, some pigments, solders, and alloys	Joints and the spine are particularly vulnerable, resulting in severe pain. It weakens bones and damages the kidneys.
CFC	Cleaning solvents, refrigerants, aerosol propellants	Risk of skin cancer and possible genetic damage.
CR	Dyes, pigments	Asthma, bronchitis, lung cancer, and damage to the liver and kidneys are possible.
Dioxins	Printed wiring boards (PWB), different types of cables, metal smelting	Increased cancer risk.
Pb	Thermoelectric elements, thermocouples, thermistors	The kidneys, reproductive system, and nervous system may be impacted. May cause blood and brain illnesses.
Li	Batteries of mobiles, photographic equipment	Long-term exposure can cause nausea, vomiting, disorientation, and muscular weakness.
Hg	Batteries, flat screen monitors, copper machines, switches	Affects the central nervous system, kidneys, immune system, and foetal development. May harm the brain and liver and cause skin issues.
PAH	Wiring, printed circuit boards	Eye discomfort, nausea, vomiting, diarrhoea, and disorientation. Long-term exposure may cause cataracts, kidney and liver damage, and jaundice.
PVC	Cables, insulation coating	Can cause respiratory and immune system damage.
PCB	Transformers, capacitors, softening agents	Damage to the immune, reproductive, neurological, and endocrine systems; also a persistent environmental pollutant.

Because many electronic components contain lead, mercury, cadmium, and brominated flame retardants, improper disposal can produce contamination that moves from local disposal sites into broader terrestrial and aquatic food webs (Table 1). For a short review, this section should stay focused on the key ecological pathways rather than listing every contaminant in detail.

3.1 Soil Contamination

When e-waste is dumped or broken down in open environments, heavy metals and additives can leach into surrounding soils. This contamination can reduce soil fertility, disturb microbial communities, and impair the activity of decomposers that support nutrient cycling and plant growth. In the long term, polluted soils may become less suitable for agriculture and less capable of sustaining diverse terrestrial ecosystems.

3.2 Water Pollution

Rainfall and surface runoff can transport soluble contaminants from e-waste sites into groundwater, streams, wetlands, and rivers. Once in aquatic systems, metals such as mercury and cadmium can accumulate in organisms and move upward through the food chain, increasing exposure at higher trophic levels. This process is especially concerning because aquatic biodiversity is sensitive to chronic, low-dose contamination that may not be immediately visible but can strongly affect reproduction, growth, and survival.

3.3 Air Pollution

Open burning of cables, plastic housings, and circuit components releases fine particles and toxic combustion products into the atmosphere. These emissions may include dioxins, furans, and other hazardous compounds that can spread beyond the original disposal site and settle on vegetation, soil, and water surfaces. Airborne contamination therefore extends the ecological footprint of e-waste well beyond the point of disposal.

3.4 Impact on Wildlife

Wildlife exposed to contaminated habitats may experience reduced fertility, developmental defects, behavioural changes, and lower survival rates. Species living near dumping grounds, informal recycling areas, or polluted waterways are particularly vulnerable because exposure can occur through food, water, sediment, and inhalation. Over time, these pressures can reduce population size and species richness, contributing to broader ecosystem instability.

3.5. Ecological Consequences

The biodiversity impact of e-waste is therefore not limited to isolated toxicity events; it reflects cumulative ecosystem stress. Soil degradation, aquatic contamination, and atmospheric release can act together, weakening ecosystem

resilience and reducing the ability of habitats to recover from disturbance. In a short review, this section can close by emphasizing that e-waste is both a waste-management issue and a biodiversity-conservation issue.

IV. E-Waste Management Practices

E-waste management is a critical step in reducing ecological damage because the environmental impact of discarded electronics depends largely on how they are collected, treated, and recovered. Effective management should prioritize safe collection, controlled dismantling, material recovery, and disposal of residues that cannot be reused or recycled. For a short review, it is useful to distinguish between formal recycling systems and informal practices, since their environmental outcomes are fundamentally different.

4.1 Formal Recycling Sector

Formal recycling refers to regulated facilities that use controlled mechanical, chemical, and thermal processes to recover metals and other useful materials while limiting pollutant release. These systems are better suited for separating valuable fractions such as copper, gold, silver, and rare earth-bearing components, while also managing hazardous residues more safely. In principle, formal recycling supports both resource recovery and pollution prevention, making it the preferred route for sustainable e-waste handling. However, formal recycling remains unevenly distributed across regions, and many countries still lack sufficient collection infrastructure, sorting capacity, and treatment facilities. As a result, a large share of e-waste is never processed through certified channels, which limits recovery rates and weakens environmental protection. Strengthening formal systems is therefore essential for reducing leakage of toxic substances into ecosystems.

4.2 Informal Recycling Sector

In contrast, informal recycling relies on manual dismantling, open burning, acid leaching, and unprotected separation of components. These methods are often used because they are cheap and accessible, but they release heavy metals, brominated compounds, and combustion by-products directly into the environment. Informal recycling therefore creates simultaneous risks for workers, nearby communities, and surrounding ecosystems. The environmental consequences are especially severe when informal processing occurs near homes, farms, or waterways. Smoke from burning cables and plastics can contaminate air and soil, while wastewater from crude chemical extraction can pollute surface water and groundwater. In biodiversity terms, these practices turn e-waste sites into localized pollution hotspots that can affect both terrestrial and aquatic species.

4.3. Management Challenges

A major challenge in e-waste management is the gap between the growing volume of discarded electronics and the limited capacity of existing recycling systems. Short product lifespans, rapid replacement cycles, and weak collection networks all contribute to this mismatch. In many regions, the result is a fragmented system in which valuable materials are lost and hazardous substances are dispersed. This challenge is not only technical but also policy-driven. Effective management requires stronger extended producer responsibility schemes, better enforcement, public collection systems, and incentives for repair and reuse. Without these measures, e-waste will continue to accumulate faster than it can be safely processed.

4.4. Transition to Sustainability

The most sustainable approach combines formal recycling with circular-economy strategies such as product repair, reuse, refurbishment, and design for disassembly. Such approaches reduce the demand for virgin raw materials and lower the amount of waste entering uncontrolled disposal streams. For biodiversity protection, the key point is that waste prevention is more effective than remediation after contamination has already occurred.

V. Case Study: Kaliganj, Nadia

Kaliganj in Nadia district, West Bengal, provides a useful micro-level example of how e-waste issues are no longer confined to major urban centers. As electronic consumption expands into smaller towns and rural areas, discarded devices increasingly enter local waste streams through household disposal, informal resale networks, and movement of waste from nearby urban regions. This makes Kaliganj relevant not only as a local case, but also as a representative example of how e-waste management challenges are spreading into semi-rural landscapes. The main concern in such areas is the absence of organized collection and recycling infrastructure. In the lack of formal systems, obsolete electronics are often stored indefinitely, dismantled by informal workers, or mixed with municipal solid waste. These practices increase the chance that toxic components will be released into soil, ponds, drainage channels, and agricultural land. For a region where ecosystems and livelihoods are closely linked, even small amounts of contamination can have disproportionate effects. A second issue is low awareness among consumers and small-scale handlers. Many residents are not fully informed about the hazards associated with batteries, circuit boards, fluorescent lamps, screens, and plastic casings containing flame retardants. As a result, devices are often discarded with household waste or sold to informal buyers without any environmental safeguards. This creates a

pathway for both resource loss and pollutant dispersal. In rural and peri-urban settings such as Kaliganj, biodiversity vulnerability is amplified by the close connection between land use, water use, and local ecology. Contaminated soil can affect crop productivity and soil organisms, while polluted water bodies may threaten fish, amphibians, birds, and other aquatic or semi-aquatic species. Because such ecosystems often have lower buffering capacity than heavily managed urban systems, contamination may persist longer and spread more easily through food webs. This case therefore highlights an important policy lesson: e-waste management must extend beyond large cities and industrial recycling hubs. Local awareness programs, community-level collection points, producer take-back systems, and safe downstream processing are needed to prevent rural contamination. In short, Kaliganj illustrates how e-waste is becoming a decentralized environmental problem, requiring local intervention as well as national regulation.

VI. Future Projections of E-Waste Generation

The volume of e-waste is expected to continue rising rapidly in the coming years because electronic consumption is increasing while product life cycles are becoming shorter. This growth is driven by several interconnected trends, including faster device replacement, expansion of connected technologies, wider access to consumer electronics, and the growing dependence of daily life on digital hardware. As a result, the environmental burden of e-waste will likely intensify unless collection and recovery systems improve at the same pace. A key factor behind this trend is planned obsolescence, both technical and behavioural. Many products are designed with limited repairability, sealed components, and short software support periods, which encourages consumers to replace devices rather than repair them. At the same time, the rapid pace of innovation creates strong market pressure for frequent upgrades, especially in smartphones, computers, entertainment devices, and household appliances. This combination increases the total quantity of discarded equipment entering waste streams each year. The future burden is not only a matter of volume but also of composition. As electronics become smaller, more complex, and more integrated, separating useful materials from hazardous constituents becomes more difficult. Devices increasingly contain multi-layered assemblies, miniature batteries, composite plastics, and finely distributed metals, which complicates recycling and raises processing costs. This means that future e-waste will likely be harder to manage than older waste streams, even if collection rates improve. In many regions, recycling infrastructure is not expanding quickly enough to keep up with this growth. Formal collection systems remain limited, and informal handling continues to dominate in many low- and middle-income settings. If this gap persists, more waste will accumulate in landfills (Figure 1), storage sites, and informal recycling centres, increasing the likelihood of contamination of soil, water, and air. From a biodiversity perspective, this is particularly concerning because the ecological effects of e-waste are cumulative and often long-lasting. Future projections therefore suggest that e-waste will remain a major environmental issue unless circular-economy approaches become mainstream. Extending product lifetimes, improving repairability, strengthening take-back systems, and redesigning electronics for disassembly can all slow the growth of waste generation. In this sense, the most effective strategy is not only better recycling, but also reducing the rate at which electronics become waste in the first place.



Figure 1. E-Waste landfilling.

VII. Strategies for Sustainable E-Waste Management

Sustainable e-waste management requires a systemic shift away from linear “produce–use–discard” models toward circular approaches that reduce waste generation and improve resource recovery. Rather than relying solely on end-of-life treatment, the most effective strategies integrate environmental considerations into every stage of the product lifecycle, from design to disposal. This not only limits the release of hazardous substances into ecosystems, but also conserves raw materials and protects biodiversity by reducing pressure on mining and land use.

7.1. Policy and Regulation

Strong regulatory frameworks are essential for channelling e-waste toward safe handling and recycling. Extended producer responsibility (EPR) schemes, in which manufacturers share responsibility for the collection and treatment of their products, have proven effective in some regions. Such policies can incentivize design for disassembly, recyclability, and the use of safer materials. Enforcement, clear liability rules, and standardized reporting are equally important to ensure that regulations translate into real-world improvements rather than mere compliance paperwork.

7.2. Recycling Infrastructure

Investing in modern, well-regulated recycling infrastructure is another critical step (Figure 2). Formal facilities should combine mechanical sorting, physical separation, and, where appropriate, controlled chemical or smelting processes to recover metals and other useful materials while minimizing environmental impact. These systems must be scaled to match local e-waste volumes and integrated with reliable collection networks so that devices do not leak into informal channels. In rural and semi-urban areas, localized, modular recycling units can help bridge the gap between generation and treatment capacity.




Figure 2. e-waste recycling process and recovery of valuable materials.

7.3. Public Awareness




Figure 3. Importance of education to create awareness about e-waste.

Awareness-raising campaigns can help households, small businesses, and informal collectors understand why proper e-waste management matters (Figure 3). Clear information on health risks, environmental impacts, and available collection points can encourage people to return devices rather than dump or dismantle them irresponsibly. Education should target schools, community centres, and local media, especially in regions where informal handling is common and environmental exposure is high.

7.4. Circular Economy Approach

Embedding circular-economy principles into electronics production and use can reduce waste at its source. Repair-support programs, second-hand markets, refurbished devices, and modular designs that allow component upgrades all extend product lifetimes. Designing for easier disassembly and material separation ensures that when devices do become waste, they can still be efficiently recycled. In this way, circularity helps both resource conservation and ecosystem protection.

7.5. Technological Innovations

Finally, technological innovation plays a key role. Advances in selective leaching, hydrometallurgical and bio-hydrometallurgical processes, and low-energy recovery methods can improve the efficiency and environmental performance of recycling. Equally important are material innovations, such as safer flame retardants, reduced use of hazardous additives, and bio-based or easily recyclable polymers. Together, these developments can reduce the toxicity burden of e-waste while supporting a more sustainable digital future.

VIII. Discussion

The link between e-waste and biodiversity is complex but increasingly clear: as electronic consumption grows, so does the release of hazardous substances into ecosystems. Soil, water, and air pollution resulting from inadequate management can disrupt nutrient cycles, alter habitat quality, and expose wildlife to chronic stress, all of which threaten species survival and ecosystem resilience. The case of Kaliganj, Nadia illustrates how these pressures are no longer limited to major cities; they are spreading to rural and semi-urban areas where ecological buffers are weaker and institutional capacity is often limited. At the same time, technological progress and circular-economy thinking offer realistic pathways for mitigation. Improved regulations, better recycling infrastructure, and stronger producer responsibility can reduce the volume of e-waste leaking into the environment while enhancing material recovery. Public awareness and education are equally important because they influence how consumers and informal handlers behave at the most local level. In many ways, the biggest challenge is not technical feasibility, but coordination: aligning corporate incentives, policy frameworks, and community practices toward a common goal. An important limitation of current knowledge is the lack of comprehensive, ecosystem-scale data on e-waste contamination gradients and their long-term biological effects. Most studies focus on specific sites or particular pollutants, which makes it difficult to generalize across regions. Future work should therefore prioritize monitoring programs that track contamination, species responses, and ecosystem functions over time. Such data would strengthen both risk assessment and policy design, helping to ensure that management strategies are proportionate to the actual ecological burden.

IX. Conclusion

E-waste represents one of the fastest-growing waste streams of the digital age, and its environmental impact extends well beyond resource depletion and pollution. When improperly managed, discarded electronics release toxic substances that can accumulate in soil, water, and food webs, endangering both human health and biodiversity. In regions like Kaliganj, Nadia, the movement of waste from urban to rural settings demonstrates how these risks are becoming more diffuse and harder to control without systemic intervention. Addressing the problem requires more than isolated cleanup efforts. It calls for integrated policies that promote circular design, support safe recycling, and encourage responsible consumption and disposal. Governments, industries, and individuals all have a role to play in reducing the toxic footprint of e-waste. By strengthening collection systems, enforcing environmental standards, and fostering innovation in materials and recycling technologies, society can move toward a more sustainable digital future in which electronic progress does not come at the cost of ecological integrity. For a short review, this manuscript has aimed to show that e-waste is not just a waste-management issue, but a biodiversity and ecosystem-protection issue. The challenge now is to translate this understanding into concrete action, so that the full life cycle of electronic devices becomes more compatible with the long-term health of the planet.

REFERENCES

- [1]. Serpe, A., Purchase, D., Bisschop, L., Chatterjee, D., De Giannis, G., Garelick, H., Kumar, A., Peijnenburg, W., Piro, V., & Cera, M. (2025). 2002–2022: 20 years of e-waste regulation in the European Union and worldwide trends in legislation and innovation technologies for a circular economy. *RSC Sustainability*, 3, 1039–1083.
- [2]. Bala, K., Dogra, A., Rajkumar, Nandini, Das, A. P., & Ali, S. (2026). Impact of e-waste on human health and environment. In *Electronic Waste and Environmental Pollution: Impacts and Sustainable Management through Circular Economy* (pp. 99–111). Springer.
- [3]. Okolo, C., Ahaneku, C., Ugwu, F., Muogbo, C., & Odoh, B. (2025). A review of the environmental and health impacts of electronic waste: Global challenges and sustainable management solutions. *International Journal of Environment and Climate Change*, 15, 220–237.
- [4]. Vaish, B., Sharma, B., Singh, P., & Singh, R. P. (2019). E-waste and their implications on the environment and human health. In *E-waste Recycling and Management: Present Scenarios and Environmental Issues* (pp. 219–232). Springer.
- [5]. Manfredi, S., Hamor, T., Wittmer, D., Nuss, P., Solar, S., Latunussa, C., Tecchio, P., Nita, V., Vidal, B., & Blengini, G. (2017). *Raw Materials Information System (RMIS): Towards v2.0*. Publications Office of the European Union.

- [6]. Smith, R. L., &Behdad, S. (2025). From present to future: A review of e-waste recycling processes. *Waste Management*, 204, 114863.
- [7]. Baldé, C., Kuehr, R., Yamamoto, T., McDonald, R., D'Angelo, E., Althaf, S., Bel, G., Deubzer, O., Fernandez-Cubillo, E., & Forti, V. (2025). Electronic waste rising five times faster than documented e-waste recycling. United Nations Report.
- [8]. Jadoun, S., Chinnam, S., Jabin, S., Upadhyay, Y., Jangid, N. K., & Zia, J. (2025). Recovery of metals from e-waste: Facts, methods, challenges, case studies, and sustainable solutions. *Environmental Science & Technology Letters*, 12, 8–24.
- [9]. Shahabuddin, M., Uddin, M. N., Chowdhury, J., Ahmed, S., Uddin, M., Mofijur, M., & Uddin, M. (2023). A review of the recent development, challenges, and opportunities of electronic waste (e-waste). *International Journal of Environmental Science and Technology*, 20, 4513–4520.
- [10]. Mohamed, D., Fayad, A., Mohamed, A.-M. O., & Al Nahyan, M. T. (2025). Transforming waste into wealth: The role of e-waste in sustainable mineral resource management.
- [11]. Jain, M., Kumar, D., Chaudhary, J., Kumar, S., Sharma, S., & Verma, A. S. (2023). Review on e-waste management and its impact on the environment and society. *Waste Management Bulletin*, 1, 34–44.
- [12]. Bagwan, W. A. (2024). Electronic waste (e-waste) generation and management scenario of India, and ARIMA forecasting of e-waste processing capacity of Maharashtra state till 2030. *Waste Management Bulletin*, 1, 41–51.
- [13]. Shoukat, A., Nurulain, S. M., Qureshi, R. N., Zafar, A., Hussain, S., Khan, S. A., Mehmood, S., Asmat, S., Raza, S., & Shah, S. T. A. (2026). Cardiometabolic dysregulation and PON1 genetic susceptibility in chronic e-waste recyclers exposed to potentially toxic elements. *Cardiovascular Toxicology*, 26, 13.
- [14]. Kumar, P., Kumar, P., & Singh Thakur, S. (2024). Waste and hazardous elements: A comprehensive study of chemical components and environmental threats. *International Journal of Innovative Science Research Technology*, 1074–1085.
- [15]. Kumar, R., Saxena, A., Chawla, J., & Kumar, V. (2024). Environmental impacts of e-waste: Pollution and resource depletion. In *Sustainable Solutions for E-waste and Development* (pp. 12–32). IGI Global Scientific Publishing.
- [16]. Hassan, K., Kaur, P., & Vijayasingam, V. (2023). E-waste—study of environmental impacts and waste management practices. In *Proceedings of the 2023 International Conference on Integrated Intelligence and Communication Systems (ICIICS)* (pp. 1–5). IEEE.
- [17]. Apprey, M. W., Dzah, C., Agbevanu, K. T., Agyapong, J. O., & Selase, G. S. (2024). E-waste management from electronic repair workshops: Societal implications and environmental consequences. *Societal Impacts*, 4, 100077.
- [18]. Oise, G., & Konyeha, S. (2025). Environmental impacts in e-waste management using deep learning. *Discover Artificial Intelligence*, 5, 210.
- [19]. Kumar, A. (2025). E-waste and the telecommunication industry: A 15-year bibliometric mapping of research progress. *Applied Research Advances*, 1, 28–49.
- [20]. Sarker, B. K., Sarker, D. K., Shaha, S. R., Saha, D., & Saha, M. K. (2025). Scenario of e-waste management: Navigating business prospects overcoming environmental challenges. *SSRN Electronic Journal*, 5261877.
- [21]. Prata, J. C. (2024). The environmental impact of e-waste microplastics: A systematic review based on the DPSIR framework. *Environments*, 11, 30.
- [22]. Aarthi, N. (2026). In-depth coverage of e-waste sources, characteristics, and environmental impact. In *Electronic Waste and Environmental Pollution: Impacts and Sustainable Management through Circular Economy* (pp. 37–52). Springer.
- [23]. Jacob, D. J. (1999). Introduction to Atmospheric Chemistry. Princeton University Press. United Nations Environment Programme (2019). Emissions Gap Report
- [24]. Forster, P. Et al. (2007). Changes in atmospheric constituents and radiative forcing. IPCC Report.
- [25]. Lelieveld, J., Crutzen, P. J., & Dentener, F. J. (1998). Changing concentration of atmospheric oxidants. *Nature*.
- [26]. Monks, P. S. Et al. (2009). Atmospheric composition change – global and regional air quality. *Atmospheric Environment*.
- [27]. Shindell, D., et al. (2012). Simultaneously mitigating near-term climate change and improving human health. *Science*.
- [28]. Solomon, S. (1999). Stratospheric ozone depletion and climate change. *Reviews of Geophysics*.
- [29]. National Oceanic and Atmospheric Administration (2021). Annual Greenhouse Gas Index. Andreae, M. O., & Crutzen, P. J. (1997). Atmospheric aerosols and climate. *Science*.
- [30]. Penner, J. E., et al. (2001). Aerosols, their direct and indirect effects. IPCC Climate Report.
- [31]. Isaksen, I. S. A., et al. (2009). Atmospheric composition change and climate–chemistry interactions. *Atmospheric Environment*.
- [32]. Ramanathan, V. Et al. (2001). Aerosols, climate, and the hydrological cycle. *Science*. Wild, M. (2009). Global dimming and brightening. *Journal of Geophysical Research*.
- [33]. Secretariat, R. S. (2011). E-waste in India. India Research Unit (LARRDIS), Rajya Sabha Secretariat, New Delhi.
- [34]. Hulgaard, T., Vehlow, J. (2010). Incineration: process and technology. *Solid Waste Technology & Management*, 1, 363–392.
- [35]. Jaiswal, A., Samuel, C., Patel, B. S., Kumar, M. (2015). Go green with WEEE: Eco-friendly approach for handling E-waste. *Procedia Computer Science*, 46, 1317–1324.
- [36]. Jaiswal, A., Samuel, C., Patel, B. S., Kumar, M. (2015). Go green with WEEE: Eco-friendly approach for handling E-waste. *Procedia Computer Science*, 46, 1317–1324.
- [37]. Jaiswal, A., Samuel, C., Patel, B. S., Kumar, M. (2015). Go green with WEEE: Eco-friendly approach for handling E-waste. *Procedia Computer Science*, 46, 1317–1324.
- [38]. Zhang, K., Schnoor, J. L., Zeng, E. Y. (2012). E-waste recycling: where does it go from here? *Environmental Science & Technology*, 46(20), 10861–10867.
- [39]. Robinson, B. H. (2009). E-waste: an assessment of global production and environmental impacts. *Science of the Total Environment*, 408(2), 183–191.
- [40]. Saha, L., Kumar, V., Tiwari, J., Rawat, S., Singh, J., Baudhdh, K. (2021). Electronic waste and their leachates impact on human health and environment: Global ecological threat and management. *Environmental Technology & Innovation*, 24, 102049.